

# Nostosin G and Spiroidesin B from the Cyanobacterium *Dolichospermum* sp. NIES-1697

Chin-Soon Phan, Jakia Jerin Mehjabin, Andrea Roxanne J. Anas, Masahiro Hayasaka, Reiko Onoki, Juting Wang, Taiki Umezawa, Kenji Washio, Masaaki Morikawa, and Tatsufumi Okino\*



Cite This: *J. Nat. Prod.* 2022, 85, 2000–2005



Read Online

ACCESS |




Metrics & More



Article Recommendations



Supporting Information



**ABSTRACT:** Chemical investigation of the cyanobacterium *Dolichospermum* sp. NIES-1697 afforded nostosin G (1), a linear tripeptide, spiroidesin B (2), and two known compounds, anabaenopeptins I (3) and J (4). Planar structures and absolute configurations for 1 and 2 were determined by 2D NMR, HRMS, Marfey's methodology, chiral-phase HPLC, and enzymatic degradation. Nostosin G (1) is a unique example of a linear peptide containing three subunits, 4-hydroxyphenyllactic acid (Hpla), homotyrosine (Hty), and argininal, with potent trypsin inhibitory properties. The biosynthetic gene clusters for nostosin G (1) and spiroidesin B (2) were investigated based on the genome sequence of *Dolichospermum* sp. NIES-1697.

The continuing exploration of natural products from cyanobacteria is a point of discussion to date. Many recent developments have been reported regarding the exploration of natural products from cyanobacteria. A few recent articles about cyanobacterial natural products are the discoveries of the cyanobactin tolypamide from *Tolypothrix* sp.,<sup>1</sup> nocuolactylates and hapalosins from *Nodularia* sp. LEGE 06071 and *Fischerella* sp. PCC9431, respectively,<sup>2</sup> cyclic lipopeptide puwainaphycins and minutissamides from *Nodularia harveyana* UHCC-0300,<sup>3</sup> pentabrominated biindole alkaloid aetokthonotoxin from *Aetokthonos hydrillicola*,<sup>4</sup> guanidine bis-prenylated peptide argicyclamide A from *Microcystis aeruginosa* NIES-88,<sup>5</sup> and chlorinated fatty acid amide columbamides F, G, and H from *Moorea bouillonii*.<sup>6</sup> Interestingly, these secondary metabolites often exhibited biological activities such as antitrypanosomal, biosurfactant, cytotoxic, and enzyme inhibitory properties.<sup>3–8</sup> In previous studies, we have isolated spumigin J from *Dolichospermum compacta* NIES-835 (formerly *Anabaena compacta*). Both spumigins A and J showed thrombin and cathepsin B inhibitory activities.<sup>7</sup> Hence, as part of our search for new bioactive natural products from cyanobacteria, we investigated an unexplored strain from the NIES collection, *Dolichospermum* sp. NIES-1697. The chemical investigation of this strain has led to the isolation of two new metabolites, nostosin G (1) and

spiroidesin B (2), and two known compounds, anabaenopeptins I (3) and J (4) (Figure 1). Herein, we report the isolation, structure elucidation, absolute configurations, trypsin inhibitory activities, and genomic analyses of biosynthetic gene clusters for nostosin G (1) and spiroidesin B (2).


## RESULTS AND DISCUSSION

Nostosin G (1) was obtained as a mixture of interconverting cyclic and linear forms (Figure S1) from the hydrophilic fraction of the *Dolichospermum* sp. NIES-1697 culture grown in 50 L of BG-11 medium. Cyclic hemiaminal formation of the argininal (Argol) residue was observed in spumigins and aeruginosins, in which the sodium borohydride (NaBH<sub>4</sub>) reduction was used to convert the mixture into a single peak containing argininal (Argol).<sup>8–11</sup> In addition to nostosin G (1), spiroidesin B (2),

Received: April 27, 2022

Published: August 10, 2022





**Figure 1.** Structures of 1–5 and chemical reduction of 1.

anabaenopeptins I (3), and J (4) were also isolated from the hydrophilic fraction.


The structure (5) of a reduced form of 1 was determined by NMR, MS, and Marfey's methodology (Table 1, Figures S16 and S17). A molecular formula of  $\text{C}_{25}\text{H}_{35}\text{N}_5\text{O}_6$  was determined based on the HRESIMS deprotonated molecule  $[\text{M} - \text{H}]^-$  at  $m/z$  500.2505. The  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra in  $\text{DMSO}-d_6$  (Table 1) showed three  $\text{sp}^3$  methines at  $\delta_{\text{C}}$  72.5, 52.1, and 50.2;  $\delta_{\text{H}}$  4.26, 4.05, and 3.72 suggested signals for three amino acid  $\text{C}\alpha$  methines, which later were confirmed to be homotyrosine (Hty), argininol (Argol), and hydroxyphenyllactic acid (Hpla). Two disubstituted benzenes were deduced from overlapped 1D NMR signals at  $\delta_{\text{C}}$  130.2, 129.0, 115.1, and 114.7;  $\delta_{\text{H}}$  7.01, 6.91, and 6.65, COSY correlations of  $\delta_{\text{H}}$  7.01 to 6.65 and 6.91 to 6.65, and HMBC correlations of signals at  $\delta_{\text{H}}$  7.01, 6.91, and 6.65 to its corresponding HSQC cross-peaks. Additional correlations from both COSY and HMBC spectra confirmed the presence of Hpla and Hty (Figure 2, Table 1). The presence of Argol was determined by the remaining correlations of both COSY and HMBC data (Figure 2, Table 1). The key HMBC peaks of the  $\text{C}\alpha$  methines at  $\delta_{\text{H}}$  4.26, 4.05, and 3.72 to carbonyl carbons at  $\delta_{\text{C}}$  173.2 and 171.0, together with selected ROESY correlations between NH signals ( $\delta_{\text{H}}$  7.79/7.84 and 7.75) and  $\text{C}\alpha$  methines ( $\delta_{\text{H}}$  4.26 and 4.05), allowed the connection of a linear peptide of Hpla-Hty-Argol (Figure 2). The presence of isomers was confirmed when most signals of Hty-Argol found in the  $^{13}\text{C}$  NMR were observed as split peaks (Figure S3). The structures of these isomers were determined as *R*-Hpla-*L*-Hty-*D*-Argol and *R*-Hpla-*L*-Hty-*L*-Argol based on Marfey's analyses of the non-reduced form and authentic Hpla chromatographic comparison (Figures S16 and S17). However, we believe nostosin G (1) was originally synthesized as *R*-Hpla-*L*-Hty-*L*-Argol due to the absence of an epimerase domain in its biosynthetic gene cluster (*nst*), as shown in Figure 3. Most analogues of spumigins and aeruginosins have four residues,<sup>7–10,12–15</sup> and only a few examples contain three units.<sup>11,16</sup>

Compound 2 has the molecular formula of  $\text{C}_{32}\text{H}_{37}\text{N}_3\text{O}_8$  with a HRESIMS deprotonated molecule  $[\text{M} - \text{H}]^-$  at  $m/z$  590.2511. The  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra in  $\text{DMSO}-d_6$  (Table

**Table 1.** NMR Spectroscopic Data ( $\text{DMSO}-d_6$ ) for 5 (Reduced Form of Nostosin G)

unit	position	$\delta_{\text{C}}$ <sup>a</sup> type	$\delta_{\text{H}}$ (J in Hz) <sup>b</sup>	HMBC <sup>c</sup>
Hpla <sup>d</sup>	1	173.2, C		
	2	72.5, CH	4.05 dd (8.6, 3.8)	1, 3, 4
	3	39.6, CH <sub>2</sub>	2.88 dt (13.8, 3.8) 2.55 ddd (13.8, 8.6, 4.1)	1, 2, 4, 5/9 1, 2, 4, 5/9
	4	128.6, C		
	5/9	130.2, CH	7.01 d (7.6)	3, 5/9, 7
	6/8	114.7, CH	6.65 m	4, 6/8
	7	155.6, C		
Hty	1	171.0, C		
	2	52.1, CH	4.26 m	1, 3, 4 Hpla 1
	3	34.9, CH <sub>2</sub>	1.85 m 1.76 m 2.39 m	1, 2, 4, 5 1, 2, 4, 5 2, 3, 5, 6/10
	4	30.3, CH <sub>2</sub>		
	5	131.3, C		
	6/10	129.0, CH	6.91 d (7.9)	4, 6/10, 8
	7/9	115.1, CH	6.65 m	5, 7/9
	8	155.3, C		
	NH		7.84 d (8.3)/7.79 d (8.3)	2, Hpla 1
	Argol	1	63.1, CH <sub>2</sub>	3.37 td (10.7, 4.8) 3.29 td (10.7, 5.8)
2		50.2, CH	3.72 m	1, Hty 1
3		27.8, CH <sub>2</sub>	1.58 m 1.31 m	2, 4, 5 2, 4, 5
4		25.1, CH <sub>2</sub>	1.49 m 1.43 m	2, 3, 5 2, 3, 5
5		40.6, CH <sub>2</sub>	3.07 m	4, 6
6		156.7, C		
NH			7.75 d (8.3)	2, Hty 1
N <sup>δ</sup> H			7.54 d (8.3)	4, 5, 6

<sup>a</sup>Measured at 150 MHz. <sup>b</sup>Measured at 600 MHz. <sup>c</sup>HMBC correlations are from the proton(s) stated to the indicated carbon. <sup>d</sup>Hpla = 4-hydroxyphenyllactic acid.




**Figure 2.** 2D NMR correlations of the reduced form of nostosin G (5) and spiroidesin B (2).

2) showed three  $C\alpha$  methines of  $\alpha$ -amino acids at  $\delta_C$  52.7, 52.3, and 51.1;  $\delta_H$  4.30, 4.26, and 4.10, three NH protons at  $\delta_H$  8.16, 8.13, and 7.91, and four carbonyl carbons at  $\delta_C$  173.5, 171.7, 171.5, and 169.6, where one carbonyl is part of an acetyl group, and the rest indicated three amino acid residues. Three sets of two methylenes and disubstituted benzenes showed the existence of three homotyrosines (Hty). The COSY and HMBC correlations (Figure 2) confirmed a linear structure of NAc-Hty-Hty-Hty. Marfey's analysis of the acid hydrolysate of 2 indicated the presence of both *L*- and *D*-Hty. The *L*- and *D*-Hty sequence was determined by the cleavage of the C-terminal amino acid of 2 using carboxypeptidase Y.<sup>17</sup> The Hty released from the reaction was purified using HPLC along with dipeptide NAc-Hty-Hty (Figure S18). The molecular formula of the dipeptide was confirmed to be  $C_{22}H_{26}N_2O_6$  from the observed  $[M + H]^+$  ion at  $m/z$  415.1876 (calculated for  $[M + H]^+$  415.1869). From Marfey's analysis of the hydrolysates of the dipeptide and the released Hty, the structure of spiroidesin B was confirmed as NAc-*D*-Hty-*D*-Hty-*L*-Hty (Figures S19–20). A close example to 2 was spiroidesin, hexanoyl-*D*-Hty-*L*-Hty-*D*-

Phe, isolated from the cyanobacterium *Anabaena spiroides*.<sup>18</sup> The known compounds anabaenopeptins I (3) and J (4) were identified in comparison with literature data.<sup>19</sup>

To reveal the biosynthetic gene clusters of these peptides, we sequenced the genome of *Dolichospermum* sp. NIES-1697. AntiSMASH software (version 5.0)<sup>20</sup> allowed us to propose the assembly lines for nostosin G (1), spiroidesin B (2), and anabaenopeptins I (3) and J (4). The putative nostosin G biosynthetic gene cluster (*nst*) was found to consist of two genes, NstA/B as nonribosomal peptide synthase (NRPS), where NstA is a ketoreducing NRPS (Figure 3). Ketoreducing NRPS modules are composed of A-KR-PCP sequences with a ketoreductase (KR) domain that catalyzed  $\beta$ -ketoreduction on  $\alpha$ -keto acid to a hydroxy acid residue.<sup>21</sup> The NRPSs were identified to be composed of three NRPS modules. They were predicted to produce a peptide containing three units consistent with the structure of nostosin G, where the NstA A<sub>1</sub> domain (VGVWIAASGK), NstB A<sub>1</sub> domain (DLGFTGCV), and NstB A<sub>2</sub> domain (DVETTGA V) activated Hpla, Hty, and Arg, respectively.<sup>14</sup> The architecture of the NRPSs was similar to a typical spumigin biosynthetic gene cluster containing four NRPS modules,<sup>9,10,14,22,23</sup> in contrast to three NRPS modules for the biogenesis of nostosin G (Figure 3). As for the biosynthesis of Hty in nostosin G, three putative enzymes (Figure 3) found in the upstream region of *nst* are believed to be involved in homotyrosine synthesis and share homology with HphA (74.74%; YP\_001865968), HphB (79.72%; YP\_001865961), and HphCD (82.24%; YP\_001865962) from homophenylalanine biosynthetic genes in cyanobacterium *Nostoc punctiforme* PCC73102.<sup>24</sup> The region upstream of *nst* encodes four putative NRPSs (Figure 3) that share homology with AptA (91.41%; WP\_006196339), AptB (85.85%; WP\_006196340), AptC (73.43%; WP\_006196341), and AptD (77.43%; WP\_006196342) from the biosynthesis of



**Figure 3.** Proposed nostosin G and spiroidesin B biosynthetic gene clusters. Domains of A, adenylation; KR, ketoreductase; T, thiolation; C, condensation; E, epimerization; TE, thioesterase; R, reductase.

**Table 2. NMR Spectroscopic Data (DMSO- $d_6$ ) for Spiroidesin B (2)**

unit	position	$\delta_C^a$ type	$\delta_H$ (J in Hz) <sup>b</sup>	HMBC <sup>c</sup>
NAc	1	169.6, C		
	2	22.5, CH <sub>3</sub>	1.89 s	1
Hty (1)	1	171.7, C		
	2	52.7, CH	4.26 td (8.6, 5.2)	1, NAc 1
	3	33.9, CH <sub>2</sub>	1.91 m	1, 2, 4, 5
			1.76 m	1, 2, 4, 5
	4	30.7, CH <sub>2</sub>	2.44–2.51 m	2, 3, 5, 6/10
	5	131.5, C		
	6/10	129.1, CH	6.95 d (8.0)	4, 6/10, 8
	7/9	115.1, CH	6.65 d (8.0)	5, 7/9
	8	155.4, C		
NH		8.16 d (8.0)	2, NAc 1	
Hty (2)	1	171.5, C		
	2	52.3, CH	4.30 td (8.6, 4.6)	1, Hty(1) 1
	3	34.5, CH <sub>2</sub>	1.92 m	1, 2, 4, 5
			1.80 m	1, 2, 4, 5
	4	30.4, CH <sub>2</sub>	2.40–2.50 m	2, 3, 5, 6/10
	5	131.4, C		
	6/10	129.1, CH	6.95 d (8.0)	4, 6/10, 8
	7/9	115.1, CH	6.65 d (8.0)	5, 7/9
	8	155.3, C		
NH		7.91 d (8.0)	1, 3, Hty(1) 1	
Hty (3)	1	173.5, C		
	2	51.1, CH	4.10 td (9.2, 4.6)	1, Hty(2) 1
	3	33.2, CH <sub>2</sub>	1.90 m	1, 2, 4, 5
			1.83 m	1, 2, 4, 5
	4	30.4, CH <sub>2</sub>	2.40–2.50 m	2, 3, 5, 6/10
	5	130.9, C		
	6/10	129.2, CH	6.91 d (8.0)	4, 6/10, 8
	7/9	115.1, CH	6.63 d (8.0)	5, 7/9
	8	155.4, C		
NH		8.13 d (8.0)	2, 3, Hty(2) 1	

<sup>a</sup>Measured at 150 MHz. <sup>b</sup>Measured at 600 MHz. <sup>c</sup>HMBC correlations are from the proton(s) stated to the indicated carbon.

anabaenopeptin in the cyanobacterium *Nodularia spumigena* CCY9414,<sup>25</sup> suggesting these four putative NRPSs are most likely associated with the biosynthesis of the isolated anabaenopeptins I (3) and J (4). This gene organization (Figure 3), where nostosin and anabaenopeptin assembly lines are co-located in the genome and are separated by a 13 kb region containing putative HphBCD, was also observed in the genome of the cyanobacterium *Sphaerospermopsis torques-reginae* ITP-024, which is also known as a producer of Hty-containing spumigins K–N.<sup>8,22</sup> A homology analysis that was carried out on the Integrated Microbial Genomes and Microbiomes (IMG/M) System using NstA as query found five homologue genes (Figure S23).<sup>26</sup> The domain structures of these homologues SpuA and SpuB shared similarities to NstA and NstB except for their genetic organization (Figure S24).

A unique feature in the gene cluster was the presence of an unrelated NRPS positioned between NstA and NstB (Figure 3). This is different from the typical *spu* assembly line (from spumigin biosynthesis), where SpuA and SpuB are located side by side, without another NRPS in between.<sup>9,10,14,22,23</sup> An attempt to predict the activated substrates for all three A domains of SprA via *in silico* prediction showed identical specificity-conferring codes DLGFTGCV, suggesting that they activated the same amino acid. We believed that this is Hty based

on the high similarity of the binding pocket sequence to the AptC A<sub>2</sub> domain (DLGFAGCV), in accordance with the structures of anabaenopeptins I (3) and J (4) found in this study strain. The binding pockets activated by the Hty residue from anabaenopeptin biosynthetic gene clusters have AptC A<sub>2</sub> domains (DLGFTGVC) in several *Planktothrix* strains and (DLGFTGCV) from *Sphaerospermopsis torques-reginae* ITP-024<sup>22,27</sup> and SpuB A<sub>1</sub> domain (DLGFTGCV) from spumigin biosynthesis (Table S1).<sup>14</sup> Thus, we reason that peptide D-Hty-D-Hty-L-Hty of spiroidesin B (2) was a product of this NRPS (SprA), spiroidesin B biosynthetic gene cluster (*spr*). The existence of epimerases found in modules 1 and 2 within SprA matched the structure determined by NMR, cleavage of the C-terminal amino acid, and Marfey's analyses. The structure of 2 has an acetyl group at the N-terminus, which is most likely catalyzed by an acetyltransferase. This can be exemplified by the MvdB acetyltransferase-catalyzed Tyr N-acetylation in the biosynthesis of microviridin K.<sup>28</sup> However, no such acetyltransferase was encoded in the regions adjacent to this NRPS. The N-acetyltransferase involved in the production of 2 may be located at a different locus; this has been seen in the N-acetylation of erythrochelin, where N-acetyltransferase EryM/Mcd (SACE\_1304) was located outside of the erythrochelin biosynthetic gene cluster.<sup>29</sup> Two putative acetyltransferases (NG743\_14015 and NG743\_14050) that share homology with the NeuD O-acetyltransferase at 95.45% (WP\_168358461.1) and 98.62% (WP\_200888902.1), respectively, were found in a different locus that was separated by a distance of 1 Mbp from the *spr* gene cluster. The acetylation likely occurs after the tethered tripeptide is hydrolytically released by the TE domain. Further study is needed to elucidate this biosynthetic pathway.

The linear peptide spumigins, aeruginosins, and nostosins are reported as serine and cysteine protease inhibitors and showed potent inhibitory activities against thrombin, trypsin, plasmin, and cathepsin B.<sup>7,12,14,16</sup> The mixture of L/D-Argal 1 (non-reduced) inhibited trypsin with an IC<sub>50</sub> value 0.1  $\mu$ M. Similar trypsin inhibitory properties were found in the related compounds, spumigin E<sup>14</sup> and nostosin A,<sup>16</sup> containing an Argal unit. The reduced form of nostosin G (5) did not show any significant activities (IC<sub>50</sub> value >200  $\mu$ M) because the Argal unit could not covalently bind with trypsin active sites.<sup>8,16</sup> Spiroidesin B (2), anabaenopeptin I (3), and anabaenopeptin J (4) were also subjected to trypsin inhibitory properties; however, none of them showed significant activities. This study discovered two new natural products, nostosin G (1) and spiroidesin B (2), from *Dolichospermum* sp. NIES-1697, whose biosynthetic gene clusters were proposed based on the genome sequence, and nostosin G (1) exhibited trypsin inhibition with an IC<sub>50</sub> value of 0.1  $\mu$ M.

## EXPERIMENTAL SECTION

**General Experimental Procedures.** Optical rotation and UV data were measured on a HORIBA SEPA-300 polarimeter and JASCO V-650 spectrophotometer, respectively, with MeOH as the solvent. The electronic circular dichroism (ECD) spectrum was taken using a JASCO J-720 spectropolarimeter. <sup>1</sup>H and <sup>13</sup>C NMR spectra were measured on a JEOL ECA 600 NMR spectrometer. Chemical shifts were denoted in  $\delta$  (ppm) relative to residual solvent peaks as internal standard (DMSO- $d_6$ ,  $\delta_H$  2.50,  $\delta_C$  39.5). High-performance liquid chromatography (HPLC) experiments were performed with a JASCO HPLC system equipped with a PU-980 pump and UV-970 UV/vis detector. HR-ESI (–) MS experiments were performed with an Agilent 1100 Series HPLC system coupled with a micrOTOF-HS mass spectrometer (Bruker Daltonics).

**Culture Conditions.** *Dolichospermum* sp. NIES-1697 was grown in BG-11 media,<sup>30</sup> with aeration (filtered air, 0.3 L/min) at 25 °C under illumination of 80  $\mu\text{E}/\text{m}^2 \text{ s}^{-1}$  on a 12L:12D cycle. Cells were collected by continuous flow centrifugation at 10 000 rpm after 4–5 weeks of incubation.

**DNA Extraction and Genome Sequencing.** High molecular weight DNA was extracted by first crushing the frozen cells with a mortar and pestle, then dissolving the powder with CTAB buffer (3% CTAB, 1.4 M NaCl, 0.2%  $\beta$ -mercaptoethanol, 20 mM EDTA, 100 mM Tris pH 8, RNase A) and incubating at 50 °C for 30 min. Later, 750  $\mu\text{L}$  of  $\text{CHCl}_3$  was added, then centrifuged with an inverted tube at 15 000 rpm for 5 min. The upper layer was gently transferred into a centrifuge tube containing 200  $\mu\text{L}$  of isopropanol. The appearance of a white string (precipitated DNA) became visible after gentle rotation of the tube; this precipitated DNA was collected for genomic sequencing using the PacBio RS II system. The 125 $\times$  coverage data were obtained. The reads were subjected to *de novo* assembly using Hierarchical Genome Assembly Process (HGAP v.2), resulting in one circular chromosome (accession number: CP099464) with 5 833 376 Mbp of sequence data, which contained 38.52% G + C and 5969 open reading frames. The genome sequence was analyzed by AntiSMASH software (version 5.0) for the detection of biosynthetic gene clusters.<sup>20</sup>

**Extraction and Isolation.** Freeze-dried cells (2.8 g from 50 L of culture in 4–5 weeks) were homogenized and extracted with MeOH (200 mL  $\times$  3). The concentrated extract was partitioned between  $\text{H}_2\text{O}$  and EtOAc. The  $\text{H}_2\text{O}$  (202.8 mg) extracts were applied to ODS (YMC-Gel, 150  $\mu\text{m}$ ) with an aqueous MeOH elution system to obtain 20%, 40%, 60%, and 100% MeOH fractions. The combined fractions of 20% and 40% MeOH fractions (100.0 mg) were purified by HPLC with a C18 column (Wakosil-II 5C18 AR, 20  $\times$  250 mm, UV detection 215 nm, flow rate 4.0 mL/min) and isocratic conditions (0–40 min, 33% MeCN with 0.1% TFA) to yield **2** (7.0 mg), **3** (1.0 mg), and **4** (7.0 mg). Reduction of a half-portion of the 60% MeOH fraction (20.0 mg from 40.0 mg) was done with  $\text{NaBH}_4$  to improve peak resolution. Subsequently, this reduction product was purified by HPLC with a C18 column (Wakosil-II 5C18 AR, 20  $\times$  250 mm, UV detection 210 nm, flow rate 4.0 mL/min) and isocratic conditions (0–20 min, 20% MeCN with 0.1% TFA) to yield **5** (8.0 mg).

**Reduced form of nostosin G (5):** colorless oil;  $[\alpha]_{\text{D}}^{27} +9.4$  ( $c$  0.67, MeOH); UV (MeOH)  $\lambda_{\text{max}}$  ( $\log \epsilon$ ) 278 (3.71), 224 (4.27) nm; ECD (0.001 M, MeOH)  $\lambda_{\text{max}}$  ( $\Delta\epsilon$ ) 226 (2.25) nm (Figure S22); NMR (DMSO- $d_6$ , 600 MHz) see Table 1; HRESIMS  $m/z$  500.2505  $[\text{M} - \text{H}]^-$  (calcd for  $\text{C}_{23}\text{H}_{34}\text{N}_5\text{O}_6$ , 500.2515).

**Spirodesin B (2):** colorless oil;  $[\alpha]_{\text{D}}^{18} -52$  ( $c$  0.4, MeOH); UV (MeOH)  $\lambda_{\text{max}}$  ( $\log \epsilon$ ) 278 (3.44), 224 (4.26) nm; NMR (DMSO- $d_6$ , 600 MHz) see Table 2; HRESIMS  $m/z$  590.2511  $[\text{M} - \text{H}]^-$  (calcd for  $\text{C}_{32}\text{H}_{36}\text{N}_3\text{O}_8$ , 590.2508).

**Oxidation, Hydrolysis, and Advanced Marfey's Analyses for the Absolute Configurations of the Amino Acids.** Nostosin G (**1**) was dissolved in 30%  $\text{H}_2\text{O}_2$ , and one drop of concentrated HCl was added. The solution was kept at room temperature for 24 h. Then the solution was lyophilized and hydrolyzed. Hydrolysis of **1** and **2** (0.2 mg) was executed by adding 200  $\mu\text{L}$  of 6 M HCl solutions. The solution was placed in a vacuum hydrolysis tube and heated at 110 °C for 16 h. The hydrolysates were dried and dissolved in 100  $\mu\text{L}$  of  $\text{H}_2\text{O}$ . The aqueous solutions of amino acid standards and hydrolysates of **1** and **2** were subjected to Marfey's analysis by adding 200  $\mu\text{L}$  of FDLA solution (in acetone) followed by 50  $\mu\text{L}$  of 1 M  $\text{NaHCO}_3$ , and the resulting solution was heated to 40 °C for 1 h.<sup>31</sup> The reaction mixture was cooled to room temperature and neutralized with 50  $\mu\text{L}$  of 2 M HCl solution. The resulting residues were resuspended in 200  $\mu\text{L}$  of MeCN and directly analyzed by LC-MS. The LC-MS analysis of arginine derivatives from Marfey's analysis was performed on a Cadenza CD-C18 column (2  $\times$  150 mm) with a linear gradient elution of 20–80% MeCN containing 0.1% formic acid for 30 min and a flow rate of 0.2 mL/min. For the analysis of homotyrosine derivatives, a linear gradient of 25% MeCN containing 0.1% formic acid to 65% MeCN containing 0.1% formic acid using a YMC-Pack Pro C18 column (2  $\times$  50 mm) and 0.2 mL/min flow rate was used. The enzymatic reaction applying carboxypeptidase Y to cleave the COOH-terminal Hty of compound **2** was done by the

previously reported method.<sup>17</sup> Compound **2** (1 mg) was dissolved in 200  $\mu\text{L}$  of 0.1 M pyridine acetate buffer (pH 5.5), and a few microliters of carboxypeptidase Y was added. The solution was incubated for 16 h and dried to remove the buffer. The remaining dipeptide and released Hty were purified using HPLC with Cosmosil MS-II (4.6  $\times$  250 mm, UV detection 215 nm, flow rate 0.5 mL/min) and gradient conditions (0–30 min, 5–30% MeCN with 0.1% TFA). The dipeptide containing two Hty moieties was hydrolyzed using 6 M HCl at 110 °C for 16 h, followed by Marfey's analysis to determine the sequence of Hty for **2**. The masses and retention time of the DLA derivatives of the L- and D-amino acid standards were as follows ( $m/z$ , L-, D-): Arg (469, 15 min, 14 min); Hty (490, 20 min, 23 min).

**Determination of the Absolute Configuration of Hydroxyphenyllactic Acid.** The acid hydrolysate of **1** was extracted with ethyl ether, which separated Hpla from the amino acid mixtures. The ether extract of the acid hydrolysates was then dried *in vacuo* and dissolved in MeOH. The MeOH solution was analyzed on an Astec Chirobiotic TAG column (250  $\times$  2.1 mm), with a flow rate of 0.3 mL/min, UV detection at 277 nm, and a linear elution with 19:1 MeOH/1% aqueous triethylammonium acetate buffer (pH 4). The coelution experiments of the Hpla derivative from **1** with an authentic standard of R/S-Hpla confirmed the absolute configuration. The retention time of the hydrolysate of **1** has a  $t_{\text{R}}$  of 3.6 min, R-Hpla has a  $t_{\text{R}}$  of 3.6 min, and S-Hpla has a  $t_{\text{R}}$  of 3.2 min.

**Trypsin Inhibition Assay.** All of the enzymes and substrates used in this study were purchased from Sigma Chemical Co. The trypsin was dissolved in 50 mM Tris-HCl (pH 7.6) to prepare 150 U/mL enzyme solutions. The  $\alpha$ -benzoyl-D/L-arginine-4-nitroanilide hydrochloride (43.3 mg) was dissolved in 1 mL of dimethyl sulfoxide (DMSO). The DMSO solution was diluted 100 times with 50 mM Tris-HCl (pH 7.6) and used as the substrate for the trypsin assay. In this study, each 200  $\mu\text{L}$  assay mixture containing 30  $\mu\text{L}$  of Tris-HCl buffer, 50  $\mu\text{L}$  of enzyme solution, and 20  $\mu\text{L}$  of test solution or inhibitor was added to each microtiter well using a 96-well plate.<sup>32</sup> The assay mixture and the substrate solution were preincubated at 37 °C for 5 min, and then 100  $\mu\text{L}$  substrate solutions were added to start the reaction. The absorbance of the well was measured immediately and after a 30 min incubation at 405 nm. Leupeptin from Peptide Institute, Inc., was used as a positive control with an  $\text{IC}_{50}$  value of 0.6  $\mu\text{M}$ .

## ■ ASSOCIATED CONTENT

### Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.jnatprod.2c00382>.

<sup>1</sup>H NMR, <sup>13</sup>C NMR, HSQC, HMBC, COSY, and ROESY spectra of compounds **1** and **2**, LC-MS chromatogram of compounds **1** and **2**, Marfey's derivatives of the hydrolysates of **1** and **2** (PDF)

## ■ AUTHOR INFORMATION

### Corresponding Author

Tatsufumi Okino – Graduate School of Environmental Science and Faculty of Environmental Earth Science, Hokkaido University, Sapporo 060-0810, Japan; [orcid.org/0000-0002-8363-0467](https://orcid.org/0000-0002-8363-0467); Email: [okino@ees.hokudai.ac.jp](mailto:okino@ees.hokudai.ac.jp); Fax: +81 11 706 4867

### Authors

Chin-Soon Phan – Faculty of Environmental Earth Science, Hokkaido University, Sapporo 060-0810, Japan;

[orcid.org/0000-0002-6500-696X](https://orcid.org/0000-0002-6500-696X)

Jakia Jerin Mehjabin – Faculty of Environmental Earth Science, Hokkaido University, Sapporo 060-0810, Japan;

[orcid.org/0000-0003-3117-7151](https://orcid.org/0000-0003-3117-7151)

**Andrea Roxanne J. Anas** – Graduate School of Environmental Science, Hokkaido University, Sapporo 060-0810, Japan; [orcid.org/0000-0002-5243-6880](https://orcid.org/0000-0002-5243-6880)

**Masahiro Hayasaka** – Graduate School of Environmental Science, Hokkaido University, Sapporo 060-0810, Japan

**Reiko Onoki** – Graduate School of Environmental Science, Hokkaido University, Sapporo 060-0810, Japan

**Juting Wang** – Graduate School of Environmental Science, Hokkaido University, Sapporo 060-0810, Japan

**Taiki Umezawa** – Graduate School of Environmental Science and Faculty of Environmental Earth Science, Hokkaido University, Sapporo 060-0810, Japan; [orcid.org/0000-0003-4280-6574](https://orcid.org/0000-0003-4280-6574)

**Kenji Washio** – Graduate School of Environmental Science and Faculty of Environmental Earth Science, Hokkaido University, Sapporo 060-0810, Japan

**Masaaki Morikawa** – Graduate School of Environmental Science and Faculty of Environmental Earth Science, Hokkaido University, Sapporo 060-0810, Japan

Complete contact information is available at:

<https://pubs.acs.org/10.1021/acs.jnatprod.2c00382>

### Author Contributions

C.-S.P. and J.J.M. contributed equally.

### Notes

The authors declare no competing financial interest.

### ACKNOWLEDGMENTS

This work was supported by JSPS KAKENHI Grant Number 19F19096. C.-S.P. is a recipient of the JSPS Postdoctoral Fellowship for Foreign Researchers (ID No. P19096).

### REFERENCES

- (1) Purushothaman, M.; Sarkar, S.; Morita, M.; Gugger, M.; Schmidt, E. W.; Morinaka, B. I. *Angew. Chem., Int. Ed.* **2021**, *60*, 8460–8465.
- (2) Figueiredo, S. A. C.; Preto, M.; Moreira, G.; Martins, T.; Abt, K.; Melo, A.; Vasconcelos, V. M.; Leão, P. N. *Angew. Chem., Int. Ed.* **2021**, *60*, 10064–10072.
- (3) Saurav, K.; Caso, A.; Urajová, P.; Hrouzek, P.; Esposito, G.; Delawská, K.; Macho, M.; Hájek, J.; Cheel, J.; Saha, S.; Divoká, P.; Arsin, S.; Sivonen, K.; Fewer, D. P.; Costantino, V. *ACS Omega* **2022**, *7*, 11818–11828.
- (4) Breinlinger, S.; Phillips, T. J.; Haram, B. N.; Mareš, J.; Martínez Yerena, J. A.; Hrouzek, P.; Sobotka, R.; Henderson, W. M.; Schmieder, P.; Williams, S. M.; Lauderdale, J. D.; Wilde, H. D.; Gerrin, W.; Kust, A.; Washington, J. W.; Wagner, C.; Geier, B.; Liebeke, M.; Enke, H.; Niedermeyer, T. H. J.; Wilde, S. B. *Science* **2021**, *371*, eaax9050.
- (5) Phan, C.-S.; Matsuda, K.; Balloo, N.; Fujita, K.; Wakimoto, T.; Okino, T. *J. Am. Chem. Soc.* **2021**, *143*, 10083–10087.
- (6) Mehjabin, J. J.; Wei, L.; Petitbois, J. G.; Umezawa, T.; Matsuda, F.; Vairappan, C. S.; Morikawa, M.; Okino, T. *J. Nat. Prod.* **2020**, *83*, 1925–1930.
- (7) Anas, A. R. J.; Kisugi, T.; Umezawa, T.; Matsuda, F.; Campitelli, M. R.; Quinn, R. J.; Okino, T. *J. Nat. Prod.* **2012**, *75*, 1546–1552.
- (8) Sanz, M.; Salinas, R. K.; Pinto, E. *J. Nat. Prod.* **2017**, *80*, 2492–2501.
- (9) Fewer, D. P.; Jokela, J.; Pauku, E.; Österholm, J.; Wahlsten, M.; Permi, P.; Aitio, O.; Rouhiainen, L.; Gomez-Saez, G. V.; Sivonen, K. *PLoS One* **2013**, *8*, No. e73618.
- (10) Jokela, J.; Heinilä, L. M. P.; Shishido, T. K.; Wahlsten, M.; Fewer, D. P.; Fiore, M. F.; Wang, H.; Haapaniemi, E.; Permi, P.; Sivonen, K. *Front. Microbiol.* **2017**, *8*, 1963.
- (11) Nowruz, B.; Wahlsten, M.; Jokela, J. *Iran. J. Biotechnol.* **2019**, *17*, 71–78.
- (12) Matsuda, H.; Okino, T.; Murakami, M.; Yamaguchi, K. *Tetrahedron* **1996**, *52*, 14501–14506.
- (13) Fujii, K.; Sivonen, K.; Adachi, K.; Noguchi, K.; Sano, H.; Hirayama, K.; Suzuki, M.; Harada, K. *Tetrahedron Lett.* **1997**, *38*, 5525–5528.
- (14) Fewer, D.; Jokela, J.; Rouhiainen, L.; Wahlsten, M.; Koskenniemi, K.; Stal, L. J.; Sivonen, K. *Mol. Microbiol.* **2009**, *73*, 924–937.
- (15) Mazur-Marzec, H.; Kaczkowska, M. J.; Blaszczyk, A.; Akcaalan, R.; Spoof, L.; Meriluoto, J. *Mar. Drugs* **2013**, *11*, 1–19.
- (16) Liu, L.; Jokela, J.; Wahlsten, M.; Nowruz, B.; Permi, P.; Zhang, Y. Z.; Xhaard, H.; Fewer, D. P.; Sivonen, K. *J. Nat. Prod.* **2014**, *77*, 1784–1790.
- (17) Hayashi, R.; Moore, S.; Stein, W. H. *J. Biol. Chem.* **1973**, *248*, 2296–2302.
- (18) Kaya, K.; Mahakhant, A.; Keovara, L.; Sano, T.; Kubo, T.; Takagi, H. *J. Nat. Prod.* **2002**, *65*, 920–921.
- (19) Murakami, M.; Suzuki, S.; Itou, Y.; Kodami, S.; Ishida, K. *J. Nat. Prod.* **2000**, *63*, 1280–1282.
- (20) Blin, K.; Shaw, S.; Steinke, K.; Villebro, R.; Ziemert, N.; Lee, S. Y.; Madema, M. H.; Weber, T. *Nucleic Acids Res.* **2019**, *47*, W81–W87.
- (21) Alonzo, D. A.; Chiche-Lapierre, C.; Tarry, M. J.; Wang, J.; Schmeing, T. M. *Nat. Chem. Biol.* **2020**, *16*, 493–496.
- (22) Lima, S. T.; Alvarenga, D. O.; Etchegaray, A.; Fewer, D. P.; Jokela, J.; Varani, A. M.; Sanz, M.; Dörr, F. A.; Pinto, E.; Sivonen, K.; Fiore, M. F. *ACS Chem. Biol.* **2017**, *12*, 769–778.
- (23) Voß, B.; Bolhuis, H.; Fewer, D. P.; Kopf, M.; Möke, F.; Haas, F.; El-Shehawry, R.; Hayes, P.; Bergman, B.; Sivonen, K.; Dittmann, E.; Scanlan, D. J.; Hagemann, M.; Stal, L. J.; Hess, W. R. *PLoS One* **2013**, *8*, No. e60224.
- (24) Koketsu, K.; Mitsuhashi, S.; Tabata, K. *Appl. Environ. Microbiol.* **2013**, *79*, 2201–2208.
- (25) Rouhiainen, L.; Jokela, J.; Fewer, D. P.; Urmann, M.; Sivonen, K. *Chem. Biol.* **2010**, *17*, 265–273.
- (26) Chen, I.-M. A.; Chu, K.; Palaniappan, K.; Ratner, A.; Huang, J.; Huntemann, M.; Hajek, P.; Ritter, S.; Varghese, N.; Seshadri, R.; Roux, S.; Woyke, T.; Eloe-Fadrosh, E. A.; Ivanova, N. N.; Kyrpides, N. C. *Nucleic Acids Res.* **2021**, *49*, D751–D763.
- (27) Christiansen, G.; Philmus, B.; Hemscheidt, T.; Kurmayer, R. *J. Bacteriol.* **2011**, *193*, 3822–3831.
- (28) Philmus, B.; Christiansen, G.; Yoshida, W. Y.; Hemscheidt, T. K. *ChemBioChem* **2008**, *9*, 3066–3073.
- (29) Lazos, O.; Tosin, M.; Slusarczyk, A. L.; Boakes, S.; Cortés, J.; Sidebottom, P. J.; Leadlay, P. F. *Chem. Biol.* **2010**, *17*, 160–173.
- (30) Waterbury, J. B.; Stanier, R. Y. *Prokaryotes* **1981**, *7*, 221–223.
- (31) Marfey, P. *Carlsberg Res. Commun.* **1984**, *49*, 591–596.
- (32) Shin, H. J.; Murakami, M.; Matsuda, H.; Yamaguchi, K. *Tetrahedron* **1996**, *52*, 8159–8168.